PCP



Street Names for PCP

Angel Dust, Animal Tranquilizer, Embalming Fluid, Ozone, Rocket Fuel, Wack, Black Dust Boat, Cliffhanger, Crystal T, Dipper, Dust joint, Goon Dust, Happy Sticks, Horse tranquilizer Kools, Lethal Weapon, Magic Dust, Magic dust, O.P.P., Paz, Peter Pan, Shermans, Trank, Wet, Ashy Larry, illy, Hog, Lovely, Ozone,.

Many of these names have been around for years, while others are modern terms that may not be as popular. Some have even gotten their roots from other countries or old stories. There is no doubt that there could be other names for this drug. Many groups especially fraternities and college-based groups, may have their own special or unique names. Familiarizing yourself with these names will empower you to make informed decisions.

Remember these can change at any time. "So Listen Up!" Our kids have learned these names and so can you.

PCP was first devolved in the 1950s as an anesthetic but due to bad side effects like: hallucinations, delirium, and mania it could no longer be approved for human medical use. It was discontinued in the 1960s by the pharmaceutical company that created it.

There are different types of PCP

PCP is illegally manufactured in labs and sold in the form of tablets, capsules, colored powders and in a liquid form. In this form the drug is known as wet.

Below are some examples of what the drug could look like.



How long does a PCP high last?

It will vary depending on how you get it in your system and it also depends on the person. The average PCP high can last anywhere from 11-52 hours.

Different doses bring different effects.

Low Dose: A low dose can give a person the feeling of euphoria, relaxation, numbness, anxiety, confusion, amnesia and blurry vision. It can also cause the person to stare aimlessly.

Medium Dose: A medium dose can cause a person to have: Confusion, Paranoia, Fever, Delusions and Schizophrenic like behavior.

High Dose: A high dose can cause the most dangerous effects. This could include: Fever, Respiratory Failure, Coma, Seizures and even Death.

Withdrawal Symptoms

Withdrawal symptoms from PCP are very dangerous and need monitoring by a medically trained professional.

These symptoms could include:

Speech difficulty, Depression, Memory loss, Weight loss, Confusion, Lack of impulse control, Anxiety, Coma, Suicide Attempts and Death.

References

http://alcoholism.about.com/od/lsd/a/pcp.htm

http://www.acadianaaddiction.com/addiction/pcp/symptoms-signs-effects

http://www.drugs.com/illicit/pcp.html

http://www.urbandictionary.com/define.php?term=PCP

http://www.drugfree.org/drug-guide/pcp/

https://faculty.washington.edu/chudler/pcp.html

http://www.isate.memphis.edu/pcp.html

http://www.isate.memphis.edu/pcp.html

https://www.youtube.com/watch?v=gag7D3CelO8

http://www.urbandictionary.com/define.php?term=Angel+Dust

Listed below are some local rehabs from the Philadelphia area

Fairmount Behavioral Health System

561 Fairthorne Avenue Philadelphia, PA 19128 215-487-4000

Website: www.fairmountbhs.com

Resources for Human Development

4700 Wissahickon Avenue, Suite #126 Philadelphia, PA 19144-4248 Office: 215-951-0300

Website: www.rhd.org